


Signs I should watch for

Raised temperature (fever) or chills

Feeling unwell

Feeling pain or discomfort in your pelvis

Redness, pain, swelling, or tenderness in your leg



Sudden shortness of breath and/or pain in your chest

Coughing up blood-streaked mucus

These symptoms indicate an **emergency**. Call the ambulance on **111** and inform them of your symptoms and recent surgery.

Contact your surgeon (and/or GP) **immediately** if you experience any of these symptoms, or any others that concern you.

Phone contacts:

Apply adhesive name label here

Your checklist to:

- Complete the pre-admission risk assessment
- Ask your doctor what blood clots treatment is needed
- Bring this brochure into hospital
- Before you go home, ensure you understand what to do, talk with your nurses and doctors and write up your recovery plan.

If applicable:

I understand my blood thinning medication plan:

- Before coming to hospital
- When I go home.

DEVELOPED BY



Blood clots and **YOU**

YOU are more at risk of developing a **BLOOD CLOT** after surgery or a procedure than when you take a long-distance flight.

What can **YOU** do to reduce your **RISK**?



Want more information?

www.southerncrosshospitals.co.nz/patient-information/blood-clots

Copyright © 2019

SCHL60 12/19



My risk factors

Blood clots can develop in your legs and pelvis and can be dangerous if they travel through your body and block blood supply to your lungs.

Admission to hospital to undertake treatment puts you at **HIGH** risk of **BLOOD CLOTS**.

If any of the following apply, **YOU** are at **HIGHER RISK** of developing a blood clot (ask your doctor what blood clots treatment is needed).

- I am having a surgical operation or procedure
- I have had a blood clot in the past
- Someone in my family has had a blood clot
- I am a smoker
- I have cancer and/or am undertaking cancer treatment or therapy
- I have a chronic illness/blood disease
- I have varicose veins
- I am over 40 years old
- I am overweight
- I am pregnant or have recently given birth
- I am on the contraceptive pill
- I am undertaking hormone replacement therapy
- I am on steroids
- I have not been physically active (e.g. walking) recently
- I have recently taken a long-distance flight

What actions can *I* take to reduce my risk of **BLOOD CLOTS**?



Drink the recommended amount of water

Adequate water intake helps blood stay fluid for good blood flow. Reduced water intake could thicken the blood.



Keep active

Frequent light activity (e.g. getting up, walking around and leg exercises) helps to move blood around the legs, reducing the risk of blood sitting and clotting in the veins.



Wear compression stockings

Compression stockings squeeze the legs, reducing the diameter of the veins and improving blood flow.

Want more information?

www.southerncrosshospitals.co.nz/patient-information/blood-clots

My recovery plan

Your doctors, nurses and the information on our website can help you to complete your plan.

My discharge date: _____

How much water should I drink everyday for the next 8 weeks?

My activity prescription

Week 1 _____

Week 2 _____

Week 3 _____

Week 4 _____

Week 5 _____

Week 6 _____

Week 7 _____

Week 8 _____

How long should I wear my compression stockings?

Medication plan:

